

#7667 MAPPING OF MICRONUTRIENTS STATUS IN SOILS UNDER MULTIVARIETAL CITRUS SINENSIS PRODUCTION FOR PRECISION AGRICULTURE

¹*Bernard N. Okafor, ²Vincent Aduramigba-Modupe and ²Olubunmi Denton

¹National Horticultural Research Institute (NIHORT), Idi-Ishin, Ibadan, Nigeria, ²Institute of Agricultural Research & Training, Obafemi Awolowo University, Moor Plantation, Ibadan, Nigeria, kpakpando2009@gmail.com

ABSTRACT

Citrus production in Nigeria is below the world average; and this is caused by poor soil management among other things. The situation is further exacerbated by blanket fertiliser application and low application of precision in soil fertility management. A study was carried out on a 34-year-old multi varietal citrus orchard under sweet orange (*Citrus sinensis*) to determine the current soil fertility status and variability of micronutrients. Soil samples were collected at a sampling depth of 0-15cm to determine soil micronutrient level. The sampling were done at 7x7m interval using a hand-held GPS while spatial analysis was done using ArcGIS software. The results showed that mean value of Mn, Fe, Cu and Zn were as follows: Mn 330.08±88.84; Fe 115.7±30.74; Cu 3.14±0.72 and Zn 8.81±3.65 mg/kg, with a kurtosis of 0.44, -0.53, -0.098 and 0.94 and skewness of 0.28, 0.43, -0.31 and -0.06 for Mn, Fe, Cu and Zn respectively. Due to the micronutrient variability of the orchard soils, it is important to use precision agriculture for management in order to improve yield and soil quality.

Keywords: mapping, micronutrients, citrus, variability, soil fertility

INTRODUCTION

Attainment of sufficiency in citrus production in Nigeria remains a challenge. This is due to limiting factors such as low availability of quality rootstock material, climate change, low agricultural input, lack of precision / poor use of fertiliser, high post-harvest losses and low value addition. Of importance in increasing citrus yield in Nigeria is the need for intensive soil fertility management as most farms in Nigeria are low in fertility due to low application/ precision in use of fertiliser and blanket fertiliser application. Nigeria currently has a fertiliser index rate of 5.5kg/ha of arable land (Worldbank, 2020). This has grossly affected soil and plant nutrient needs leading to micronutrient deficiency and crop failure in many cases. Consequently, there is need for improved micronutrient fertility in Nigerian agriculture. Micronutrients are essential in crop growth and are usually required in little quantity. Raja (2009) observed that the role of micronutrient in crop production include improved crop quality, increase in yield, improved disease resistance and prevention of physiological disorder. Beyond the supply of micronutrients, soil organic matter needs to be given serious attention as there is a strong correlation between organic matter and most crop properties (Ogunkunle and Awotoye, 2011).

Precision in agricultural practice has improved its productivity significantly (Yousefi and Radzari (2014). Continuous cultivation and low application of fertiliser has led to soil fertility decline (Ogunjimi *et al.*, 2017). Due to the inherent variability of soil properties as a result of land use (Borusiewicz, 2016) and climatic change (Pareek, 2017), it is important to use precise management to ensure soil conservation and improve its quality. Precision agriculture has been a viable tool for soil variability management (Zude-Sasse *et al.*, 2016) as

it helps to reduce low and poor utilization of farm input due to variation in factor of agricultural production. Shaato *et al.*, (2005) observed that studies on micronutrient in Nigerian soil are limited. Therefore, this study was carried out to map micronutrient status of a citrus orchard in order to enhance precision in its fertility management.

MATERIALS AND METHODS

Soil samples were collected from multi-varietal citrus collection block at National Horticultural Research Institute (NIHORT), Ibadan. It consists of 12 sweet orange varieties planted at a spacing of 7m x 7m. The varieties include Bende, Meran, Umudike, Agege, Pineapple and Parson Brown among others. Soil samples were taken from 0 - 15 cm depth using a soil auger. The samples were analyzed for micronutrients using standard laboratory procedures. Data were subjected to descriptive statistics while spatial analysis was carried out using GIS software package.

RESULTS AND DISCUSSION

Table 1 shows the descriptive statistics of the micronutrient concentration of the study area. Micronutrients play a vital role in citrus productivity. As shown in Table 1, the mean values of Mn, Fe and Zn are adequate for citrus production while mean concentration of Cu was indicative of low nutrient concentration for citrus production as recommended by (Chen *et al.*, 2007). Improved fertility management of soil micronutrient is important because of its effect on crop quality and productivity (Patil *et al.*, 2018). Coefficient of variation (CV) is an indicator of the level of variability of soil properties. Wilding and Dress (1983), characterized CV values as follows; 0-15 (low variability), 15-35 (moderate variability) and >35 (high variability). From the foregoing, (Mn, Fe and Cu) were moderately variable while Cu was highly variable. Consequently, Cu fertility management should be more precise and site specific. Application of fertiliser without recourse to site specific needs of the soil/crop leads to wastage of resource and poor crop yield (schellberg *et al.*, 2008). Precision agriculture helps to restrict resource use to particular area, improve yield and farm efficiency; thereby reducing wastage (Chan, 2006). Mapping aids visibility and presentation of soil properties to enhance farm and soil management. The micronutrient maps of the citrus orchard are presented in Figures 1-2

Table 1. Descriptive statistics of the orchard.

Micronutrient	Mean	CV	Kurtosis	Skewness
Mn	330.08±88.84	29.91	0.44	0.28
Fe	115.7±30.74	26.56	-0.53	0.43
Cu	3.14±0.72	22.85	-0.01	-0.31
Zn	8.81±3.65	36.83	0.94	-0.06




Figure 1. Distribution of Cu within the orchard.’




Figure 2. Distribution of Zn within the orchard.

Mean value of Cu (3.14 mg/kg) across the orchard indicated low nutrient status but the map (Fig 1) showed particular areas of deficiency. Precision and site-specific management are required to improve Cu concentration of the orchard, enhance yield, improve soil quality and reduce wastage of fertiliser. Blanket fertiliser application should be avoided.

REFERENCES

- Borusiewicz A, Kapela K, Drożyner P, and Marczuk T. 2016. Application of Precision Agriculture Technology in Podlaskie Voivodeship. *Agricultural Engineering* 20: 5-11.
- Chan CW. 2006. Application of precision agriculture technologies in the tropical greenhouse environment. *Acta Horti*. 710: 479-484D. OI:10.17660/ActaHortic. 2006.710.59 DOI: 10.17660/ActaHortic.2006.710.59.
- Chen F, Lu JW, Liu D. 2007. Investigation of soil fertility in citrus orchards of Southern China. *Better Crops* 91 3: 24 -27.
- Ogunkunle CO, Awotoye O. 2011. Soil Fertility Status under Different Tree Cropping System in a Southwestern Zone of Nigeria. *Notulae Scientia Biologicae* 3(2): 123-128 DOI: 10.15835/nsb.3.2.6050
- Pareek N. 2017. Climate change impact on soils: adaptation and mitigation. *MOJ Eco Environ Sci* ;2(3):136-139. DOI: 10.15406/mojes.2017.02.00026
- Patil KD, More SS, Wahane MR, Puranik UY, Khobragade NH. 2018. Micronutrient importance in high tech horticulture. *Journal of Pharamcognosy and photochemistry* 7(3): 628-635.
- Raja MD. 2009. Importance of micronutrient in the changing horticultural scenario in India. *Journal of Horticultural science* 4(1): 1-27.
- Schellberg J, Hill MJ, Gerhards R, Rothmund M, Braun M. 2008. Precision agriculture on grassland: Applications, perspectives and constraints. *European Journal of Agronomy* 29(2-3): 59-71
- Shaato R, Ajayi SO, Ojanuga AG. 2005. Total and extractable copper, iron, manganese and zinc in major agricultural soils in the Lower Benue Valley, Central Nigeria and the concept of extractant efficiency. *Nigerian Journal of Chemical Research* 59 – 68.
- Wilding LP, Dress LR. 1983. Spatial variability and pedology. In: Wilding LP, Smeeck NE and Hall GF. Eds. *Pedogenesis and soil taxonomy*. Pudoc wageningen.
- Yousefi1 MR, Razdari AM. 2016. Application of GIS and GPS in Precision Agriculture (A Review). *International journal of Advanced Biological and Biomedical Research* 4(2): 473-476.
- Zude-Sasse M, Fountas S, Gemtos TA, Abu-Khalaf N. 2016. Applications of precision agriculture in horticultural crops. *Eur. J. Hortic. Sci.* 81(2): 78–90 DOI: 10.17660/eJHS.2016/81.2.2